



Tuberculosis (TB) risk assessment worksheet CY2021

This model worksheet should be considered for use in performing TB risk assessments for health-care facilities and nontraditional facility-based settings. Facilities with more than one type of setting will need to apply this table to each setting.

Scoring ✓ or Y = Yes	X or N = No	NA = Not Applicable
-----------------------------	-------------	---------------------

1. Incidence of TB

<p>What is the incidence of TB in your community (county or region served by the health-care setting), and how does it compare with the state and national average? What is the incidence of TB in your facility and specific settings and how do those rates compare? (Incidence is the number of TB cases in your community the previous year. A rate of TB cases per 10,000 persons should be obtained for comparison.) * This information can be obtained from the state or local health department.</p>	<p>Broward County</p> <p>Community rate: Decrease 2.2 (2022) 2.3 (2021) Dropped from 48 to 44 incidences with the population for Broward County of 1,963,035; 2022 incidence rate rank of 21; 2022 Case Rank 3.</p> <p>State rate: Increase 2022 incidence rate (per 100,000) 2.4 (2022) 2.29 (2021)</p> <p>National rate: 2.5 (2022) 2.37 (2021)</p> <p>Facility rate: CY 2022 Increase 2.4</p> <p>(# of confirmed diagnosed cases of TB/number of admissions)</p> <p>3/12,434 = 2.4 per 10,000 admissions 2022</p> <p>2/12,501 = 1.6 per 10,000 admissions 2021</p> <p>3/ 11,642 = 2.6 per 10,000 admissions 2020</p> <p>7/ 13,542 = 5.2 per 10,000 admissions 2019</p>
--	--

Are patients with suspected or confirmed TB disease encountered in your setting (inpatient and outpatient)?	Yes												
If yes, how many patients with suspected and confirmed TB disease are treated in your healthcare setting in 1 year (inpatient and outpatient)? Review laboratory data, infection-control records, and databases containing discharge diagnoses.	<table border="1"> <thead> <tr> <th></th> <th>Suspected</th> <th>Confirmed</th> </tr> </thead> <tbody> <tr> <td>2022:</td> <td>32</td> <td>3</td> </tr> <tr> <td>2021:</td> <td>12</td> <td>2</td> </tr> <tr> <td>2020:</td> <td>29</td> <td>3</td> </tr> </tbody> </table>		Suspected	Confirmed	2022:	32	3	2021:	12	2	2020:	29	3
	Suspected	Confirmed											
2022:	32	3											
2021:	12	2											
2020:	29	3											
Currently, does your health-care setting have a cluster of persons with confirmed TB disease that might be a result of ongoing transmission of <i>Mycobacterium tuberculosis</i> within your setting (inpatient and outpatient)?	No												

2. Risk Classification

Inpatient settings									
How many inpatient beds are in your inpatient setting?	409 (379 + 30 IRU)								
How many patients with MTB disease are encountered in the inpatient setting in 1 year? Review laboratory data, infection-control records, and databases containing discharge diagnoses.	<table border="1"> <tbody> <tr> <td>CY 2022</td> <td>3</td> </tr> <tr> <td>CY 2021</td> <td>2</td> </tr> <tr> <td>CY 2020</td> <td>3</td> </tr> <tr> <td>CY 2019</td> <td>7</td> </tr> </tbody> </table>	CY 2022	3	CY 2021	2	CY 2020	3	CY 2019	7
CY 2022	3								
CY 2021	2								
CY 2020	3								
CY 2019	7								
Depending on the number of beds and TB patients encountered in 1 year, what is the risk classification for your inpatient setting (≥ 200 beds)? (See Appendix C.) According to the CDC guidelines 2005, a “low risk” facility has less than 6 TB patients a year. A “medium risk” facility has greater than or equal to 6 confirmed cases of tuberculosis annually.	For 2022, there were 32 possible cases of MTB, with 3 cases confirmed positive, 3 pulmonary and 0 cases of extra-pulmonary infection. Our risk classification is “ low risk ”.								
Does your health-care setting have a plan for the triage of patients with suspected or confirmed TB disease?	Yes								

3. Screening of HCWs for *M. tuberculosis* Infection

Does the health-care setting have a TB screening program for HCWs?	Yes
<p>If yes, which HCWs are included in the TB screening program? (Check all that apply.)</p> <ul style="list-style-type: none"> ✓ Physicians ✓ Mid-level practitioners (nurse practitioners [NP] and physician’s assistants [PA]) ✓ Nurses ✓ Administrators ✓ Laboratory workers ✓ Respiratory therapists ✓ Physical therapists Contract staff (Required by the contracting department. Records kept in contracting department) Construction or renovation workers (same as contract workers) ✓ Service workers 	<ul style="list-style-type: none"> ✓ Janitorial staff ✓ Maintenance or engineering staff ✓ Transportation staff ✓ Dietary staff ✓ Receptionists ✓ Trainees and students (Medical students-under GME; Nursing, Respiratory and other Allied dept. under Learning/Nursing department. Records and compliance are managed by the above departments) ✓ Volunteers <p>o Others: Pharmacy, Radiology</p>
Is baseline skin testing performed with two-step TST (Tuberculin Skin Test) for HCWs?	Yes

Is baseline testing performed with QFT (QuantiFERON) or other BAMT (Blood Assay for Mycobacterium Tuberculosis) for HCWs?	No, not for 2022
How frequently are HCWs tested for <i>M. tuberculosis</i> infection?	Annually during their anniversary hire period.
Are the <i>M. tuberculosis</i> infection test records maintained for HCWs?	Yes
Where are the <i>M. tuberculosis</i> infection test records for HCWs maintained? Who maintains the records?	Employee Health Department
If the setting has a serial TB screening program for HCWs to test for <i>M. tuberculosis</i> infection, what are the conversion rates for the previous years?† (2022): 0.3% (2021): 0.0% (2020): 0.250% (2019): (raw: 19) (2018): 0.2% (2017): 0.7%	
Has the test conversion rate for <i>M. tuberculosis</i> infection been increasing or decreasing, or has it remained the same over the previous 5 years? (Check one)	Increased by 0.05% from 2020 and 0.3% from 2021. Staying within the same range.
Do any areas of the health-care setting (e.g., waiting rooms or clinics) or any group of HCWs (e.g., lab workers, emergency department staff, respiratory therapists, and HCWs who attend bronchoscopies) have a test conversion rate for <i>M. tuberculosis</i> infection that exceeds the health-care setting's annual average?	No. While not above the annual average, there were 4 conversions this year that represent an increase from the previous year. Reached target of lower than 1%. None were involved in exposure at the hospital. All worked in different departments including non-clinical.
For HCWs who have positive test results for <i>M. tuberculosis</i> infection and who leave employment at the health setting, are efforts made to communicate test results and recommend follow-up of latent TB infection (LTBI) treatment with the local health department or their primary physician?	Yes. New hire positive skin test results are screened with a chest x-ray and are referred to their PCP or community resource for evaluation of latent TB status. This is required by day 30 after first day of employment. Employees who converted are seen by an ID physician through workers comp. If employees are terminated before they are seen and evaluated, a letter is sent by employee health to follow up with workers comp, private primary care physician or their new employee health department. Exposure follow up for employees who were terminated before the 10 th week of follow up are notified by letter to follow up with their PCP or new employee health department.

4. TB Infection-Control Program

Does the health-care setting have a written TB infection-control plan?	Yes – in the Infection Prevention Plan and a Broward Health policy
Who is responsible for the infection-control program?	Chairman of Infection Prevention Committee.
When was the TB infection-control plan first written?	01/1994
When was the TB infection-control plan last reviewed or updated?	3/2021
Does the written infection-control plan need to be updated based on the timing of the previous update (i.e., >1 year, changing TB epidemiology of the community or setting, the occurrence of a TB outbreak, change in state or local TB policy, or other factors related to a change in risk for transmission of <i>M. tuberculosis</i>)?	Yes

Does the health-care setting have an infection-control committee (or another committee with infection control responsibilities)?	Yes
If yes, which groups are represented on the infection-control committee? (Check all that apply.)	<ul style="list-style-type: none"> ✓ Laboratory personnel ✓ Health and safety staff ✓ Administrator ✓ Risk assessment ✓ Quality control (QC) ✓ Environmental staff ✓ Respiratory ✓ Facilities management
<ul style="list-style-type: none"> ✓ Physicians ✓ Nurses ✓ Epidemiologists ✓ Engineers ✓ Pharmacists ✓ Nutritional staff 	

5. Implementation of TB Infection-Control Plan Based on Review by Infection-Control Committee

Has a person been designated to be responsible for implementing an infection-control plan in your health-care setting? If yes, list the name:	Yes. Dr. Indulekha Gopal, Infection Control Committee Chairman
Based on review of the medical records, what is the average number of days for the following:	
<ul style="list-style-type: none"> • Presentation of patient until collection of specimens 1 • Specimen collection until receipt by laboratory 1 • Receipt of specimen by laboratory until smear results are provided to healthcare provider 1 • Diagnosis until initiation of standard antituberculosis treatment 1 • Receipt of specimen by laboratory until culture results are provide for healthcare provider 1 • Receipt of drug susceptibility results until adjustment of antituberculosis treatment, if indicated (can take up to a few weeks) • Admission of patient to hospital until placement in airborne infection isolation (AII) 1 	
Through what means (e.g., review of TST or BAMT conversion rates, patient medical records, and time analysis) are lapses in infection control recognized?	Review of laboratory results, outbreak investigations and other means of surveillance.
What mechanisms are in place to correct lapses in infection control?	Process improvements, outbreak investigation, literature search, multidisciplinary teamwork, reporting through committee process within the facility.
Based on measurement in routine QC (Quality Control) exercises, is the infection-control plan being properly implemented?	Yes
Is ongoing training and education regarding TB infection-control practices provided for HCWs?	Yes

6. Laboratory Processing of TB-Related Specimens, Tests, and Results Based on Laboratory Review

Which of the following tests are either conducted in-house at your health-care setting's laboratory or sent out to a reference laboratory?	In-house	Sent out
Acid-fast bacilli (AFB) smears	✓	
Culture using liquid media (e.g., Bactec and MB-BacT)	✓	
Culture using solid media	✓	
Drug-susceptibility testing		✓
Nucleic acid amplification (NAA) testing		✓
Does the laboratory at your healthcare setting or the reference laboratory used by your healthcare setting report AFB smear results for all patients within 24 hours of receipt of specimen? What is the procedure for weekends?	Yes. The same process is utilized on nights and weekends as regular business hours. The laboratory will page the on call Epidemiologist to communicate positive AFB	

	results outside of normal business hours. _____
--	--

7. Environmental Controls

Which environmental controls are in place in your health-care setting? (Check all that apply and describe)

Environmental control

- ✓ All rooms (airborne infection isolation rooms)
- ✓ Local exhaust ventilation (enclosing devices and exterior devices)
- ✓ General ventilation (e.g., single-pass system, recirculation system.)
- ✓ Air-cleaning methods (e.g., high-efficiency particulate air [HEPA] filtration and ultraviolet germicidal irradiation [UVGI])

What are the actual air changes per hour (ACH) and design for various rooms in the setting?

Operating Rooms: 20 ACH
 AII Rooms: 12 ACH
 Cath Lab: 15 ACH
 Bronchoscopy Room (in GI suite): 12 ACH
 Interventional Radiology Procedure Room - 15 ACH

Which of the following local exterior or enclosing devices such as exhaust ventilation devices are used in your health-care setting? (Check all that apply)

- ✓ Laboratory hoods
- ✓ Booths for sputum induction

What general ventilation systems are used in your health-care setting? (Check all that apply)

- ✓ Single-pass system
- ✓ Constant air volume (CAV)
- ✓ Recirculation system

What air-cleaning methods are used in your health-care setting? (Check all that apply)

HEPA filtration

- ✓ Fixed room-air recirculation systems

How many AII rooms are in the health-care setting?

MICU room #1
 PACU Room # 16
 Rm: 363
 RM:618
 Rm: 620
 Rm: 622
 Rm: 820
 Rm: 822
 Rm:824
 Rm: 828
 Rm:916
 Rm: 918
 Rm:920
 Rm:922
 TICU Room #8
 ENDO Room 3 (Bronch Suite)
 ED Green Pod Room 3
 ED Orange Pod Room 46
 ED Purple Pod Room 31
 ED Yellow Pod Room 15
 B Side Room 19 (to be eliminated)
 C Side Room 26 (to be eliminated)

What ventilation methods are used for AII rooms? (Check all that apply) <u>Primary (general ventilation):</u> ✓ Single pass heating, ventilating, and air conditioning (HVAC) ✓ Recirculating HVAC systems <u>Secondary (methods to increase equivalent ACH):</u> ✓ Fixed room recirculating units. ✓ HEPA Filtration	
Does your healthcare setting employ, have access to, or collaborate with an environmental engineer (e.g., professional engineer) or other professional with appropriate expertise (e.g., certified industrial hygienist) for consultation on design specifications, installation, maintenance, and evaluation of environmental controls?	Yes
Are environmental controls regularly checked and maintained with results recorded in maintenance logs?	Yes
Are AII rooms checked daily for negative pressure when in use?	Yes
Is the directional airflow in AII rooms checked daily when in use with smoke tubes or visual checks?	Yes
Are these results readily available?	Yes
What procedures are in place if the AII room pressure is not negative?	Patient is transferred. Facilities are notified and the room is closed until pressure is confirmed negative.
Do AII rooms meet the recommended pressure differential of 0.01-inch water column negative to surrounding structures?	Yes

8. Respiratory-Protection Program

Does your health-care setting have a written respiratory-protection program?	Yes												
Which HCWs are included in the respiratory protection program? (Check all that apply) ✓ Physicians ✓ Mid-level practitioners (NPs and PAs) ✓ Nurses ✓ Administrators ✓ Laboratory personnel Contract staff Construction or renovation staff ✓ Service personnel ✓ Janitorial staff ✓ Maintenance or engineering staff ✓ Transportation staff ✓ Dietary staff Students													
Are respirators used in this setting for HCWs working with TB patients? If yes, include manufacturer, model, and specific application (e.g., ABC model 1234 for bronchoscopy and DEF model 5678 for routine contact with infectious TB patients). <table border="0" style="width: 100%; border-collapse: collapse;"> <tr> <td style="text-align: center;"><u>Manufacturer</u></td> <td style="text-align: center;"><u>Model</u></td> <td style="text-align: center;"><u>Specific application</u></td> </tr> <tr> <td>3M corporation</td> <td>N-95#1860 & 1860S</td> <td>Routine Contact with Infectious TB patients</td> </tr> <tr> <td>ProGear N95 Respirator</td> <td>RP88020 (Regular)</td> <td>Routine Contact with Infectious TB patients</td> </tr> <tr> <td>ProGear N95 Respirator</td> <td>RP88010 (Small)</td> <td>Routine Contact with Infectious TB patients</td> </tr> </table>	<u>Manufacturer</u>	<u>Model</u>	<u>Specific application</u>	3M corporation	N-95#1860 & 1860S	Routine Contact with Infectious TB patients	ProGear N95 Respirator	RP88020 (Regular)	Routine Contact with Infectious TB patients	ProGear N95 Respirator	RP88010 (Small)	Routine Contact with Infectious TB patients	
<u>Manufacturer</u>	<u>Model</u>	<u>Specific application</u>											
3M corporation	N-95#1860 & 1860S	Routine Contact with Infectious TB patients											
ProGear N95 Respirator	RP88020 (Regular)	Routine Contact with Infectious TB patients											
ProGear N95 Respirator	RP88010 (Small)	Routine Contact with Infectious TB patients											
Is annual respiratory-protection training for HCWs performed by a person with advanced training in respiratory protection?	Yes												
Does your health-care setting provide initial fit testing for HCWs? If yes, when is it conducted? ___ On hire by employee health _____	Yes												
Does your health-care setting provide periodic fit testing for HCWs? If yes, when, and how frequently is it conducted? Yearly	Yes												

What method of fit testing is used? Describe. Hood/Taste ____1. Fit check: Saccharin or Bitrex fit check. Individual is asked to do normal, deep breathing; bend over; side to side and up/down head movements). _____ _____	
Is qualitative fit testing used?	Yes
Is quantitative fit testing used? (Available)	No

9. Reassessment of TB risk

How frequently is the TB risk assessment conducted or updated in the health-care setting?	Yearly
When was the last TB risk assessment conducted?	03/2022
What problems were identified during the previous TB risk assessment? No problems were identified.	
What actions were taken to address the problems identified during the previous TB risk assessment? Not applicable.	
Did the risk classification need to be revised because of the last TB risk assessment?	No. Our risk remained the same.
Recommendations: 1. Continue annual PPD testing and/or symptom screening and x-ray review of all employees and volunteers. 2. Continue to closely monitor all patients admitted for suspected/known TB for appropriate isolation practices. 3. Continue referring new employees for latent TB infection evaluation as indicated. 4. Continue education on a yearly basis and as needed. 5. Retrak concurrent monitoring of compliance with mandatory requirement (including PPD testing). Entry to the facility is restricted until all mandatory requirements are fulfilled.	

- * If the population served by the health-care facility is not representative of the community in which the facility is located, an alternate comparison population might be appropriate.
- † Test conversion rate is calculated by dividing the number of conversions among HCWs by the number of HCWs who were tested and had prior negative results during a certain period (see Supplement, Surveillance and Detection of *M. tuberculosis* infections in Health-Care Settings).